

# Mossbauer Effect Measurement of Intermetallic Compounds in Iron-Tin System : Fe<sub>5</sub>Sn<sub>3</sub> and FeSn

著者	YAMAMOTO Hisao
journal or publication title	Science reports of the Research Institutes, Tohoku University. Ser. A, Physics, chemistry and metallurgy
volume	19
page range	213-213
year	1967
URL	<a href="http://hdl.handle.net/10097/27377">http://hdl.handle.net/10097/27377</a>

## Mössbauer Effect Measurement of Intermetallic Compounds in Iron-Tin System: $\text{Fe}_5\text{Sn}_3$ and $\text{FeSn}$ \*

Hisao YAMAMOTO

*The Research Institute for Iron, Steel and Other Metals*

### Abstract

Hyperfine fields for an Fe atom in intermetallic compounds,  $\text{Fe}_5\text{Sn}_3$  and  $\text{FeSn}$ , have been investigated by using the Mössbauer effect. The hyperfine field of  $\text{Fe}_5\text{Sn}_3$  with hexagonal  $\text{B8}_2$ -type structure is 187kOe on 2(a) sites while that on 2(d) sites has two values, 247kOe and 208kOe, at liquid-air temperature. The  $\text{B8}_2$ -type structure of  $\text{Fe}_5\text{Sn}_3$  is stable above  $760^\circ\text{C}$  and the magnetic Curie temperature is  $(280^\circ \pm 5^\circ)\text{C}$ .  $\text{FeSn}$  of hexagonal  $\text{B35}$ -type is antiferromagnetic with the Néel temperature at  $97^\circ\text{C}$  and the hyperfine field is 163kOe on an Fe site at liquid-air temperature.

Temperature dependence of hyperfine field, isomer shift and quadrupole interaction was also studied.

---

\* The **1327th** report of the Research Institute for Iron, Steel and Other Metals. Published in the Journal of the Physical Society of Japan, **21** (1966), 1058.